

references

Return to [Questions & Answers](#)

Fossett, N. G., Byrne, B.J., Kelley, S.J., Tucker, A.B., Arbour-Reily, P. and Lee, W.R. (1994). The influence of large deletions on the mutation frequency induced by tritiated water and X-radiation in male *Drosophila melanogaster* post- meiotic germ cells. *Mut. Res.* 307(1): 213-222.

Johnson, J.R., Myers, D.K., Jackson, J.S., Dunford, D.W., Gragtmans, N.J., Wyatt, H.M., Jones, A.R. and Percy, D.H. (1995). Relative biological effectiveness of tritium for induction of myeloid leukemia in CBA/H mice. *Radiat Res* 144(1):82-89.

Prestwick, W.V. and Kwok, C.S. (1993). The quality factor for the degradation spectrum of ^3H in water. *Rad. Prot. Dosimetry* 46(3): 195-200.

Straume, T and Carsten, A.L. (1993). Tritium radiobiology and relative biological effectiveness. *Health Phys.* 65: 657-672.

Tanaka, K., Sawada, S. and Kamada, N. (1994). Relative biological effectiveness and dose rate effect of tritiated water on chromosomes in human lymphocytes and bone marrow cells. *Mut. Res. Letters* 323(1-2): 53-61.

Return to [Questions & Answers](#)